Draft version February 13, 2009
Preprint typeset using LaTeX style emulateapj v. 10/09/06

arXiv:0809.2786v2 [astro-ph] 13 Feb 2009

REVISITING THE EFFECT OF H_2, HD AND LIH MOLECULES IN THE COOLING OF PRIMORDIAL GAS.

Joaquin P. Prieto[1], Leopoldo Infante[1], Raul Jimenez[2]

Draft version February 13, 2009

ABSTRACT

We use a non-equilibrium chemical network to revisit and study the effect of H_2, HD and LiH molecular cooling on a primordial element of gas. We paid special attention in the variation of HD abundance. We solve both the thermal and chemical equations for a gas element with an initial temperature $T = 1000K$ and a gas number density in the range $n_{tot} = 1 - 10^3 cm^{-3}$. These are typical propierties of the first halos which formed stars. At low densities, $n_{tot} < 10^2 cm^{-3}$, the gas reaches temperatures $\sim 100K$ and the main coolant is H_2, but at higher densities, $n_{tot} > 10^2 cm^{-3}$, the HD molecule dominates the gas temperature evolution and the gas reaches temperatures well below $100K$. The effect of LiH is negligible in all cases. We studied the effect of HD abundance on the gas cooling. The HD abundance was set initially to be in the range $n_{HD}/n_H = 10^{-7} - 10^{-5}$. The simulations show that at $n_{tot} > 10^2 cm^{-3}$ the HD cooling dominates the temperature evolution for HD abundances greater than $10^{-6} n_H$. This number decrease at higher densities. Furthermore, we studied the effect of electrons and ionized particules on the gas temperature. We followed the gas temperature evolution with $n_{H_+}/n_H = 10^{-4} - 10^{-2}$. The gas temperature reached lower values at high ionization degree because electrons, H^+ and D^+ are catalizers in the formation paths of the H_2 and HD molecules. Finaly, we studied the effect of an OB star, with $T_{eff} = 4 \times 10^4 K$, would have on gas cooling. It is very difficult for a gas with n_{tot} in the range between $1 - 100 cm^{-3}$ to drop its temperature if the star is at a distance less than $100pc$.

Subject headings: cosmology: early universe - cosmology: theory - galaxies: intergalactic medium - atomic processes

1. INTRODUCTION

In a ΛCDM Universe, the first luminous objetcs were formed due to fall-in of gas inside the dark matter potential wells (for a review see Barkana & Loeb (2001)). In order to let the fall-in of gas inside the dark matter potential wells, the gas thermal energy should have been radiated away by some physical mechanism. Since the primordial molecular clouds would have zero metallicity, the collisional excitation of the existing molecules is the most plausible mechanism for the cooling of baryonic matter in this environment because the collisional excitation cooling of both H and He atoms are inefficient at temperatures lower than $8000K$, which is higher than the temperature of star formation clouds.

Tegmark et al. (1997) showed that the first lumious objects may have formed at $z \sim 30$ inside a $10^6 M_\odot$ halo ($T_{vir} \sim 1000K$), which have recently been confirmed by O'Shea & Norman (2007) and Gao et al. (2007). Tegmark et al. (1997) also showed that the stars formed in this environment could be as massives as $\sim 100 M_\odot$ (Abel et al. 2002).

After the recombination era, the most abundant molecule in the Universe is molecular hydrogen H_2. Despite of its low primordial abundance, $\sim 10^{-3} - 10^{-4} n_H$ (Palla et al. 1983), this molecule has a fundamental role in the gas cooling at temperatures less than $8000K$. The first authors to highlight the role of H_2 in this context were Saslaw & Zipoy (1967) and Peebles & Dicke (1968). Saslaw & Zipoy (1967) showed the importance of the charge transfer reaction between H_2^+ and H to form H_2, and Peebles & Dicke (1968) suggested a mechanism to form H_2 from H^-.

The H_2 molecule forms by the H^- and H_2^+ channels mainly. The reactions

$$H + H^+ \rightarrow H_2^+ + \gamma \qquad (1)$$

$$H + e \rightarrow H^- + \gamma \qquad (2)$$

are followed by

$$H_2^+ + H \rightarrow H_2 + H^+. \qquad (3)$$

$$H^- + H \rightarrow H_2 + e. \qquad (4)$$

Due to its zero dipolar moment only quadrupolar rotational transitions are allowed, $J \rightarrow J \pm 2$, where J is the quantum number for angular momentum; Abgrall et al. (1982). Furthermore, due to its small moment of inertia (the smallest one between all molecules) the energy gap between its rotational quantum states, ΔE, is large compared with other molecules ($\Delta E_{J \rightarrow J \pm 2} \propto 1/I$, where I is the moment of inertia). The smallest energy gap is $\Delta E_{2 \rightarrow 0} \approx 500K$. With this energy difference it is very diffucult to reach temperatures below $\sim 100K$, (see Palla (1999) and references therein).

The HD molecule forms through D^+ and D channel mainly (see Dalgarno et al. (1973); Galli & Palla (2002))

$$D^+ + H_2 \rightarrow H^+ + HD, \qquad (5)$$

$$D + H_2 \rightarrow H + HD. \qquad (6)$$

---

[1] Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Santiago, Chile; jpprieto@astro.puc.cl, linfante@astro.puc.cl

[2] ICREA & Institute of Space Sciences (CSIC-IEEC), Campus UAB, Bellaterra 08193, Spain; raulj@astro.princeton.edu

HD has a moment of inertia greater than the H_2 one. Furthermore, it has a finite dipolar moment which allows internal dipolar trasitions (transitions of the kind $J \rightarrow J \pm 1$) and internal transition rates greater than in H_2, Abgrall et al. (1982). Due to its small moment of inertia, the differences between its energy states are smaller than the energy differences of the H_2 molecule. All these properties make the HD molecule an efficient cooler at low temperatures, below $\sim 100K$, (see Nagakura & Omukai (2005); Ripamonti (2007); McGreer & Bryan (2008); Palla (1999) and references therein).
The LiH molecule is formed mainly by radiative association of Li and H and associative detachment of Li^- and H (Stancil et al. 1996):

$$Li + H \rightarrow LiH + \gamma. \tag{7}$$

$$Li^- + H \rightarrow LiH + e^-. \tag{8}$$

Moreover, this molecule have both a dipolar moment and a moment of inertia larger than the ones of the HD molecule. These characteristics could make the LiH molecule an efficient cooler at low temperatures, but the cooling functions depend on the number density of the specie, so if the abundance of LiH is too low as expected in primordial environments (Stancil et al. 1996) its cooling effect will be negligible. For a review of Li chemestry see Bodo (2001) and Bodo et al. (2003).

In the work of Galli & Palla (1998) the effect of HD and LiH was included on the gas cooling. To calculate the photo-destruction ratescoefficient, for photoionization, photodetachment, and photodissociation, they assumed detailed balance with CMB photons. But, to study the effect of the first stars on the primordial gas we need the cross section for each photo-destruction process, in the spirit of Glover & Jappsen (2007). These cross sections are described below. Our current work includes the photo-destruction cross sections of both Li and Li^- and the photodisociation of LiH (in its rovibrational ground state) in contrast to previous work.

We improve over recent works (Glover & Jappsen 2007; Glover & Abel 2008) that have studied primordial cooling by exploring the effect of the HD abundance and the inclusion of a stellar radiation field. This paper is organized as follow. In §2 we describe both the thermal and chemical model required to follow the evolution of the gas temperature. In §3 we present results and discussion. It includes the gas temperature evolution as a function of gas density and molecular coolers; the gas temperature evolution as a function of HD abundance; the temperature evolution as a function of the ionization degree and finaly we show the effect of a star radiation field on the gas temperature. In §4 we present the conclusions.

2. THERMAL AND CHEMICAL MODEL.

As argued above, in a realistic cooling model of primordial gas, it is mandatory to include the molecular coolers. The main molecular coolers at low temperatures are H_2, HD and LiH. In adition to the previous molecules, the model should include the main species created in the primordial nuclesynthesis. Our model includes 21 species: H, H^+, H^-, H_2, H_2^+, H_3^+, He, He^+, He^{++}, HeH^+, D, D^+, HD, HD^+, H_2D^+, Li, Li^+, Li^-, LiH, LiH^+ and e^-. The reactions considered for these species are described in table 1, 2 and 3. This table does not include the formation of H_2 by three body reaction because this reaction is not relevant at densities studied in this work. The cooling processes (see table 3) considered in this work are:

- Collisional ionization: the gas loses energy by ionization of the different species in the environment.
- Recombination: the gas loses energy by free electrons capture.
- Collisional excitation: the gas loses energy when collisional excited electrons move to the unexcited states.
- Bremsstrahlung: loss of energy due to the radiation emited by accelerated electrons.
- Compton cooling: interchange of energy between free electrons and CMB radiation.

The molecular cooling functions are constructed as in Galli & Palla (1998). For both H_2 and HD molecules we adopted the approximate density dependent relation for the cooling functions. For LiH we adopted the low density limit cooling function. The molecular cooling functions are trated as in Puy et al. (1993) at temperatures near T_{CMB}.
The first stars, by definition, formed in an environment without previous star. But, once population III stars are formed they can photoionize the halos where new stars will form. This process is quantified by the frequency dependent cross section, $\sigma_A(\nu)$, of the reaction $A^i + \gamma \rightarrow A^{i+1} + e^-$, where the specie A^i in the i ionization state moves to $i+1$ ionization state due to the interaction with photons.

The presence of a radiation field triggers a photo-destruction rate:

$$IR_A = 4\pi \int_{\nu_{th}}^{\infty} \sigma_A(\nu) \frac{i(\nu)}{h\nu} d\nu, \tag{9}$$

where n_A is the number density of specie A; $i(\nu)$ is the specific intensity of radiation in the environment and h is Planck's constant. The integral is calculated from the threshold frequency of ionization, ν_{th}, to infinity.

The heating of the gas due to photoionizations (heat in $erg/cm^3 s$) is given by

$$\Gamma_A = n_A 4\pi \int_{\nu_{th}}^{\infty} \sigma_A(\nu) \frac{i(\nu)}{h\nu} (h\nu - h\nu_{th}) d\nu, \tag{10}$$

where $h\nu_{th}$ is the threshold energy of ionization. Strictly, in the last two expressions $i(\nu)$ should be multiplied by $1 - e^{-\tau}$, where $\tau = \int \sigma_A(\nu) n_A dl$ is the optycal depth. Here we assume $\tau >> 1$. For example, H photoionization cross section is $\sim 10^{-18} cm^2$, τ is greater than 1 for $dl > 1 - 10^{-4} pc$. If we take n_H in the range $1 - 10^4 cm^{-3}$. This distance is well below the halo scale distance.
The cross sections, the ionization rates and the heatings included in this work are mentioned in table 5
With each one of these functions we can calculate both the abundance of all species and the temperature evolution. The abundance of specie A in the gas evolves following the equation

$$\frac{\partial n_A}{\partial t} = C_{A,ij} - n_A D_{l,ph}, \tag{11}$$

where $C_{A,ij} = \Sigma_{ij} n_i n_j k_{RC,ij}$ is the creation rate of specie A by i and j with $k_{RC,ij}$ the rate coefficient of the reaction and $D_{l,ph}$ is the destruction rate of specie A by specie l, $D_l = \Sigma_l n_l k_{RC,Al}$ or by photoionization $D_{ph} = k_{IR}$. These couppled deferential equations are solved by the backward differencing formula, BDF (Aninos et al. 1997) The gas temperature T changes according to

$$\frac{\partial T}{\partial t} = \frac{\gamma - 1}{k_B \sum_i n_i}(\Gamma - \Lambda), \qquad (12)$$

where γ is the adiabatic gas index; k_B is Boltzmann's constant; Γ is the photoionization heating and Λ is the cooling by the processess mentioned before. In the range of both densities and temperatures studied here the gas can cool by molecular de-excitation. This is valid, i.e, for H_2 molecule until densities grater than $n \sim 10^8/x_{H_2} cm^{-3}$, with x_{H_2} the H_2 fraction, at higher densities the molecule reach the LTE.

3. RESULTS AND DISCUSION.

We follow the gas temperature for different densities, different HD abundances, different ionization conditions and different radiation conditions. The abundances relatives to H at the begining of the evolution are (from Galli & Palla (1998) at $z \approx 10 - 20$): $n_{H^+} = 10^{-4}$, $n_{H^-} = 10^{-12}$,$H_2^+ = 10^{-14}$, $n_{H_2} = 10^{-3}$, $n_{H_3^+} = 10^{-18}$, $n_D = 4 \times 10^{-5}$, $n_{D^+} = 0$, $n_{HD} = 10^{-6}$, $n_{HD^+} = 10^{-18}$, $n_{H_2D^+} = 10^{-19}$, $n_{He^+} = He^{++} = 0$, $n_{HeH^+} = 10^{-12}$, $n_{Li} = Li^+ = 10^{-10}$, $n_{Li^-} = 10^{-23}$, $n_{LiH} = 10^{-21}$ and $n_{LiH^+} = 10^{-18}$. The H and He densities are $\rho_H = 0.75\rho_{tot}$ and $\rho_{He} = 0.24\rho_{tot}$, where ρ_{tot} is the total baryonic matter.



FIG. 1.— Temperature evolution for an isolated element of gas with the initial abundances mentioned in the text. There are four diferent values of H density: long dashed line $n = 1cm^{-3}$; short dashed line $n = 10cm^{-3}$; dotted line $n = 100cm^{-3}$; solid line $n = 1000cm^{-3}$. Here, different colors take into account different molecular coolers: black line are H_2 coolings; red lines are H_2 and HD coolings, and H_2, HD and LiH coolings. The diference between the curves with LiH and without LiH is neglegible.

Figure 1 shows the temperature evolution of an isolated gas element with initial temperature of $1000K$. There are four different values for H abundance. These four cases show that the temperature has a strong gas density dependence; a ten times more dense gas element cools almost ten times faster in this range of density. The LiH cooling seems insignificant at this temperatures and densities, whereas the effect of HD cooling seems to be important at low temperatures when the elements of gas are more dense. In spite of the high HD cooling efficiency, the gas element can not drop its temperature below the CMB temperature (Johnson & Bromm (2006); Yoshida et al. (2007)). CMB temperature, $T_{CMB} = 2.73(1 + z)$, at $z = 20$ is $T_{CMB} \approx 57K$. This suggests the IMF of pop III stars not depend on the gas metallicity only but on the redshift z at which the stars are formed, too. For example, if we take the T_{CMB} as the lowest temperature reached by the gas, $M_J(20) \approx 0.5 \times M_J(30)$, with $M_J(z)$ the Jeans mass at redshift z.

In order to know which cooling effect dominates the evolution of gas temperature we considered the main cooling processess independently through their thermal evolution. Figure 2 shows that the cooling of H_2, HD, LiH -LiH do not appear due to its low value- and the total cooling, Λ, divided by $n_H n_{H_2}$ for a gas element with $n_{tot} = 100cm^{-3}$ and an H_2 abundance of $n_{H_2}/n_H = 10^{-3}$.

$$\frac{\Lambda}{n_H n_{H_2}} \approx \lambda_{H_2} + \left(\frac{n_{HD}}{n_{H_2}}\right)\lambda_{HD} + \left(\frac{n_{LiH}}{n_{H_2}}\right)\lambda_{LiH}. \qquad (13)$$

The last expresion correspond to the total molecular cooling. Here $\Lambda_A = n_H n_A \lambda_A$ is the cooling due to the excitation of molecule A by an H atom.

This figure shows clearly that in a gas with a number density greater than $n_{tot} \approx 10^2 cm^{-3}$ the HD cooling is similar to H_2 cooling at a temperature $\sim 100K$ when n_{HD} is greater than $10^{-6} n_H$. Then, with number densities greater than $\sim 10^2 cm^{-3}$ the HD cooling dominates the thermal evolution and the gas temperature can reach values much lower than the values reached in a gas with lower densities. This behaviour depends strongly on HD abundance, which was taken in the range $10^{-5} - 10^{-7}$ relative to H. At high densities the gas needs less HD abundance to drop its temperature. Puy & Signore (1998) found that at high densities an temperatures $\sim 200K$ the main molecular cooler is HD. This result was confirmed by Omukai (2000), Uehara & Inutsuka (2000), Flower et al. (2000), Nagakura & Omukai (2005) and Ripamonti (2007) and in a simple way with the results of figure 1 and 2. On the other hand, in order to be an effecient cooler LiH should increase its abundance in about ten order of magnitude due to its low abundance. This increment is too large to be real. The LiH cooling is not important in these cases (see Mizusawa et al. (2005)).

Figure 3 shows the gas temperature evolution of an ionized gas. This figure shows that a gas with a large ionization degree could reach lower temperatures than a gas with a small ionization degree, see for example Nagakura & Omukai (2005). This is possible because the formation of the molecular coolers, H_2 and HD, need H^+, D^+ and free electrons. The temperature difference between the $n_{H^+}/n_H = 10^{-4}$ and the $n_{H^+}/n_H = 10^{-2}$



FIG. 2.— H_2 and HD at low temperatures. In panel (a) and (b) the gas number density is $n_{tot} = 100 - 1000 cm^{-3}$, respectively. H_2 cooling in solid line; HD cooling in dotted line. The diferent colors indicate diferent initial HD abundances: $n_{HD}/n_H = 10^{-5}$ in black; $n_{HD}/n_H = 10^{-6}$ in red; $n_{HD}/n_H = 10^{-7}$ in green.

case is $\sim 100K$ at $100cm^{-3}$.

Figure 4 shows the effect of an OB star ($R_* = 15R_\odot$ and $T_{\rm eff} = 4 \times 10^4$) radiation field on the gas temperature at different distances. This figure shows that is very difficult for an element of gas to drop its temperature if there is an OB star at a distance less than $\sim 100pc$ due to photo-destruction of its cooling molecular agents, Yoshida et al. (2007). In other words if inside a halo of $\sim 10^6 M_\odot$ a star is born, it is very difficult for its surounded gas reach the necesary conditions to form other stars (Oh & Haiman (2003); Omukai & Nishi (1999)). The formation of more than one star could be possible if the seed clumps of gas evolve at the same time, without radiation feedback between them.



FIG. 3.— Gas temperature as a function of the ionization degree of a gas element with $n_{tot} = 100cm^{-3}$. In solid line $n_{H+}/n_H = 10^{-2}$; long dashed $n_{H+}/n_H = 10^{-4}$.



FIG. 4.— Temperature evolution for a gas radiated by an OB star at different distances: $50pc$ in blue; $100pc$ in black; 10^3pc in red. There are three diferent values of gas density: solid line $n_{tot} = 1cm^{-3}$; dotted line $n_{tot} = 10cm^{-3}$; dashed line $n_{tot} = 100cm^{-3}$.

The case in which the star is at $1000pc$ and the gas has $n_{tot} = 100cm^{-3}$ present the same evolution of an isolated element of gas.

4. CONCLUSIONS.

In this work we have developed a model for the temperature evolution of a primordial gas including 21 different chemical species including reaction rates and cross sections available in the literature. We have paid careful attention to explore the space parameter in abundance, specially HD, and to include the LiH to study in detail at what abundances the molecular coolants are relevant.

The main results are the following:

1. The HD molecule dominates the gas cooling at temperatures below $\sim 100-200K$ in the range of densities $10^2-10^3 cm^{-3}$ for HD abudances over $10^{-6} n_H$. The HD effect is more evident at higher densities.
2. The LiH molecule does not have a clear effect on the gas cooling. The gas would need an abundance at least ten orders of magnitude higher to be an efficient cooler, so LiH is ruled out as an important cooler in primordial gas (Mizusawa et al. (2005)).
3. A gas with high ionization degree can drop its temperature more than a neutral gas because both the ionized H and D and electrons are catalizers in the formation of H_2 and HD cooling molcules. These ionization conditions could be presents in post-shock waves zones or relic HII regions, Johnson & Bromm (2006).
4. Is very difficult for a gas to drop its temperature in the presence of an OB star located closer than $100pc$. So, in order to form more than one star in a primordial halo the formation of seed clumps should almost be instantaneous, otherwise the radiation feedbak of first stars will suppress the star formation conditions (Omukai & Nishi (1999)).

This work, as previos ones, suggests the importance of studing the effect firts stars would have on their sorrounding gas in the formation of more than one star within primordial halos (see e.g. Jimenez & Haiman (2006)). For a more accurate study, we need to follow the star formation in a hydrodinamical model, which will we present in forthcoming papers.

We are grateful to CONICYT, MECESUP and FONDAP grants. The work of RJ is supported by grants from the Spanish Ministry for Science and Innovation and the European Union (FP7 program).

REFERENCES

Abel T., Anninos P., Yu Zhang, Norman N. L., 1997, New Astr. 2, 181.

Abel, T., Bryan, G. L., & Norman, M. L. 2002, Science, 295, 93

Abgrall H., Roueff E., Viala Y., 1982, A&A Sup. 50, 505.

Adams N. G., Smith D., 1981, ApJ 248, 373.

Adams N. G., Smith D., 1985, ApJ 294, L63.

Aldrovandi S. M. V., & Pequignot D., 1973, A&A 25, 137.

AMDIS database 1989.

Anninos P., Zhang Yu, Abel T., Norman M. L., 1997, New Astr. 2, 209.

Barkana R. & Loeb A., 2001, Phys. Rep. 349, 125.

Black J. H., 1978, ApJ 222, 125.

Black J. H., 1981, MNRAS 197, 553.

Bodo E., 2001, Chem. Phys., 271, 309.

Bodo, E., Gianturco F. A., Martinazzo, R., 2003, Phys. Reports, 384, 85.

Bougleux E. Galli D., 1997, MNRAS 288, 638.

Dalgarno A., & Lepp S., 1987, in: Astrochemestry, eds. M. S. Vardya & S. P. Tarafdar, Reidel, Dordrecht, p. 109.

Dalgarno A., Black J. H. & Weisheit J. C., 1973, Astrophys. Lett., 14, 77.

Dalgarno A., Kirby K., Stancil P. C., 1996, ApJ 458, 397.

Datz S., Larsson M., Stromhölm C. et al., 1995, Phys. Rev. A. 52, 2901.

de Jong T., 1972, A&A, 20, 263.

Dunn G. H., 1968, Phys. Rev., 172, 1.

Flower D. R., Le Bourlot J. Pineau des Forêst G., Roueff E., 2000, MNRAS 314, 753.

Flower D. R. & Harris G. J., 2007, MNRAS, 377, 705.

Galli D., Palla F., 1998, A&A 335, 403.

Galli D., Palla F., 2002, Planet. Space Sci., 50, 1197.

Gao, L., Yoshida N., Abel T., Frenk C. S., Jenkins A. & Springel V., 2007, MNRAS, 314, 753.

Gerlich D., 1982 in Symposium on Atomic and Surface Physic, ed. W. Lindenger, F. Howorka & T. D. Mark (Dordrecht: Kluwer) 304.

Gerlich D., Horning S., 1992, Chem. Rev. 92, 1509.

Glover S. C. O. & Jappsen A. K., 2007, ApJ 666, 1.

Glover, S. C. O., & Abel, T. 2008, MNRAS, 388, 1627

Herbst E., 1982, A&A 111, 76.

Janev R. K., Langer W. D., Evans K. Jr., & Post D. E., 1987, Elementary Processes n hydrogen-helium plasma, Springer, Berlin.

Jimenez, R., & Haiman, Z. 2006, Nature, 440, 501

Johnson J. L. & Bromm V., 2006, MNRAS, 366, 247.

Karpas Z., Anicich V., Huntress W. T., 1979, J. Chem. Phys. 70, 2877.

Kimura M., Lane N. F., Dalgarno A., Dixon R.G., 1993, ApJ 405, 801

Kimura M., Dutta C., Shimakura N., 1994, ApJ 430, 435.

Kirby K. & Dalgarno A., 1978, ApJ, 224, 444.

Larsson M., Lepp, Dalgarno A. et al., 1996, A&A 309, L1.

Launay J. M., Le Douneuf M., Zeippen C. J., 1991, A&A 252, 842.

Lepp S., Shull J. M., 1984, ApJ 280, 465.

McGreer I. D. & Bryan G. L., 2008, arXiv0802.3918

Mac Low M. & Shull J. M., 1986, ApJ, 302, 585.

Mielke S. L., Lynch G. C., Truhlar D. G. & Schwenke D. W., 1994, J. Phys. Chem., 98, 8000.

Millar T. J., Bennet A., Herbst E., 1989, ApJ 340, 906.

Mizusawa H., Omukai K., & Nishi R., 2005, PASJ, 57, 951.

Nagakura T. & Omukai K., 2005, MNRAS, 364, 1378.

Nakamura F. & Umemura M., 2002, ApJ 569, 549.

O'Neil S. V. & Reinhardt W. P., 1978, J. Chem. Phys., 69, 2126.

O'Shea B. W. & Norman M. L. 2007, 654, 66.

Oh, S. P., & Haiman, Z. 2003, MNRAS, 346, 456.

Omukai K. & Nishi R., 1999, ApJ, 518, 64.

Omukai K., 2000, ApJ 534, 809.

Orient O. J., 1977, Chem. Phys. Lett. 52, 264.

Osterbrock D. E., 1974, Astrophysics of Gaseous Nebulae, Freeman & Co, San Francisco.

Osterbrock D. E., 1989, Astrophysics of Gaseous nebualae and active Galactic Nuclei (Mill Valley University science books).

Palla F., Salpeter E. E. & Stahler S. W., 1983, ApJ 271, 632.

Palla F., Star Formation 1999, Proceedings of Star Formation 1999, held in Nagoya, Japan, June 21 - 25, 1999, Editor: T. Nakamoto, Nobeyama Radio Observatory, p. 6-11

Peart B., Hayton D. A., 1994, J. Phys. B 27, 2551.

Peebles P. J. E., Dicke R. H., 1968, ApJ 154, 891.

Peebles P. J. E., 1971, Physical Cosmology, Princeton Univercity Press, NJ.

Peterson J. A., Alberth W. H., Moseley J. T., Sheridan J. R., 1971, Phys. Rev. A 3, 1651.

Poulaert G., Brouillard F.,Claeys W., McGowan J. W., Van Wassenhave G., 1978, J. Phys. B. 11, L671.

Puy, D., Alecian, G., Le Bourlot, J., Leorat, J., & Pineau Des Forets, G. 1993, A&A, 267, 337.

Puy D., Signore M., 1997, New Ast. 2, 299.

Puy D., Signore M., 1998b, New Ast. 3, 247.

Ramsbottom C. A., Bell K. L., Berrington K. A., 1994, J. Phys. B 27, 2905.

Reilman R. F., Manson F. T., 1979, ApJS 40, 815.

Ripamonti E., 2007, MNRAS 376, 709.

Roberge W., Dalgarno A., 1982, ApJ 255, 489.

Saslaw W. C., Zipoy D., 1967, Nature 216, 967.

Savin D. W., 2002, ApJ, 566, 599.

Savin D. W., Krstic P. S., Haiman Z. & Stancil P. C., 2004, ApJ, 606, L167.

Schneider I. F., Dulieu O., Giusti-Suzor A., Roueff E., 1994, ApJ 424, 983.

Schulz G. J., Asundi R. K., 1967, Phys. Rev., 158, 25.

Shapiro P. R. & Kang H., 1987, ApJ, 318, 32.

Shavitt I., 1959, J. Chem. Phys. 31, 1359.

Sidhu K. S., Miller S., Tennyson J., 1992, A&A 255, 453.

Stancil P. C., Babb J. F., Dalgarno A., 1993, ApJ 414, 672.

Stancil P. C., Lepp S., Dalgarno A., 1996, ApJ 458, 401.

Stancil P. C., Zigelman B., 1996, ApJ 472, 102.

Steigman G., Romano D. & Tosi M., 2007, MNRAS, 378, 576.

Stibbe D. T. & Tennyson J., 1999, ApJ, 513, L147.

Stromhölm C., Schneider I. F., Sundström G. et al., 1995, Phys. Rev. A 52, R4320.

Sundström G., Mowat J. R., Danared H. et al. 1994, Science 263, 785.
Tegmark M., Silk J., Rees M. J., Blanchard, A., Abel T. & Palla F., 1997, ApJ, 474, 1.
Theard L. P., Huntress W. T., 1974, J. Chem. Phys. 60, 2840.
Tytler D., Fan, X. M. & Burles S., 1996, Nature, 381, 207.
Uehara H. & Inutsuka S., 2000, ApJ 531, L91.
Vasiliev E. O. & Shchekinov Yu A., 2005, astro-ph/0507603v1
Verner D. A., Ferland G. J., 1996, ApJS 103, 467.
Wishar A. W. 1979, MNRAS, 187, 59P.
Yan M., Sadehpour H. R. & Dalgarno A., 1998, ApJ, 496, 1044.
Yoshida N., Omukai K. & Hernquist L., 2007, ApJ, 667, L117.
Yousif F. B., Mitchell J. B. A., 1989, Phys. Rev. A 40, 4318.
Zigelman B., Dalgarno A., Kimura M., Lane N. F., 1989, Phys. Rev. A, 40, 2340.

TABLE 1
Rate coefficents for chemical reactions.

Reaction rate (cm^3/s)	reaction	ref
Reactions with H		
RC1A97	$H + e \rightarrow H^+ + 2e$	39
RC2A97	$H^+ + e \rightarrow H + \gamma$	1
RC3GP98	$H + e \rightarrow H^- + \gamma$	40
RC4GP98	$H + H^- \rightarrow H_2 + e$	18
RC5GP98	$H + H^+ \rightarrow H_2^+ + \gamma$	10
RC6GP98	$H_2^+ + H \rightarrow H_2 + H^+$	15
RC7GP98	$H_2 + H^+ \rightarrow H_2^+ + H$	29
RC8GP98	$H_2 + e \rightarrow 2H + e$	35
RC9A97	$H_2 + H \rightarrow 3H$	20
RC10A97	$H^- + e \rightarrow H + 2e$	39
RC11A97	$H^- + H \rightarrow 2H + e$	1
RC12GP98	$H^- + H^+ \rightarrow 2H$	25
RC13GP98	$H^- + H^+ \rightarrow H_2^+ + e$	26
RC14GP98	$H_2^+ + e \rightarrow 2H$	30
RC15A97	$H_2^+ + H^- \rightarrow H_2 + H$	7
RC16GP98	$H_2^+ + H_2 \rightarrow H_3^+ + H$	38
RC17GP98	$H_2 + e \rightarrow H^- + H$	31
RC18GP98	$H_3^+ + H \rightarrow H_2^+ + H_2$	33
RC19GP98	$H_3^+ + e \rightarrow H_2 + H$	37
RC20GP98	$H_2 + H^+ \rightarrow H_3^+ + \gamma$	12
RC74GA08	$H + H + H \rightarrow H_2 + H$	9
RC74Inv	$H_2 + H \rightarrow H + H + H$	9
RC75GA08	$H + H + H_2 \rightarrow H_2 + H_2$	9
RC75Inv	$H_2 + H_2 \rightarrow H + H + H_2$	9
Reactions with He		
RC21A98	$He + e \rightarrow He^+ + 2e$	14
RC22GP98r		38
RC22A97d	$He^+ + e \rightarrow He + \gamma$	4
RC23A97	$He^+ + e \rightarrow He^{++} + 2e$	5
RC24GP98	$He^{++} + e \rightarrow He^+ + \gamma$	39
RC25GP98	$He + H^+ \rightarrow He^+ + H$	16
RC26GP98	$He^+ + H \rightarrow He + H^+$	42
RC27GP98	$He + H^+ \rightarrow HeH^+ + \gamma$	27
RC28GP98	$He + H^+ \rightarrow HeH^+ + \gamma$	27, 16
RC29GP98	$He + H_2^+ \rightarrow HeH^+ + H$	6
RC30GP98	$He^+ + H \rightarrow HeH^+ + \gamma$	42
RC31GP98	$HeH^+ + H \rightarrow He + H_2^+$	15
RC32GP98	$HeH^+ + e \rightarrow He + H$	41
RC33GP98	$HeH^+ + H_2 \rightarrow H_3^+ + He$	23

References. — (1) Abel et al. (1997); (2) Adams & Smith (1981); (3) Adams & Smith (1985); (4) Aldrovandi & Pequignot (1973); (5) AMDIS (1989); (6) Black (1979); (7) Dalgarno & Lepp (1987); (8) Datz et al. (1995); (9) Flower & Harris (2007) ;(10) Galli & Palla (1998); (11) Gerlich (1982); (12) Gerlich & Horning (1992); (13) Herbst (1982); (14) Janev et al. (1987); (15) Karpas et al. (1979); (16) Kimura et al. (1993); (17) Larsson et al. (1996); (18) Launay et al. (1991); (19) Lepp & Shull (1984); (20) MacLow & Shull (1986); (21) Mielke et al. (1994); (22) Millar et al. (1989); (23) Orient (1977); (24) Peart & Hayton (1994). (25) Peterson et al. (1971); (26) Poulaert et al. (1978); (27) Roberge & Dalgarno (1982); (28) Savin (2002); (29) Savin et al. (2004); (30) Schneider et al. (1994); (31) Schulz & Asundi (1967); (32) Shavitt (1959); (33) Sidhu et al. (1992); (34) Stancil et al. (1993); (35) Stibbe & Tennyson (1999); (36) Stromholm et al. (1995); (37) Sundstrom et al. (1994); (38) Theard & Huntress (1974); (39) Verner & Ferland (1996); (40) Wishar (1979); (41) Yousif & Mitchell (1989); (42) Zigelman et al. (1989).

TABLE 2
Rate coefficents for chemical reactions.

Reaction rate (cm^3/s)	reaction	ref
Reactions with D		
RC34A97	$D + e \rightarrow D^+ + 2e$	14
RC35GP98	$D^+ + e \rightarrow D + \gamma$	1
RC36GP98	$D + H^+ \rightarrow D^+ + H$	28
RC37GP98	$D^+ + H \rightarrow D + H^+$	28
RC38GP98	$D + H \rightarrow HD + \gamma$	19
RC39GP98	$D + H_2 \rightarrow H + HD$	21
RC40GP98	$HD^+ + H \rightarrow H^+ + HD$	15
RC41GP98	$D^+ + H_2 \rightarrow H^+ + HD$	11
RC42GP98	$HD + H \rightarrow H_2 + D$	32
RC43GP98	$HD + H^+ \rightarrow H_2 + D^+$	11
RC44GP98	$HD + H_3^+ \rightarrow H_2 + H_2D^+$	3, 22
RC45GP98	$D + H^+ \rightarrow HD^+ + \gamma$	10
RC46GP98	$D^+ + H \rightarrow HD^+ + \gamma$	10, 34
RC47GP98	$HD^+ + e \rightarrow H + D$	36
RC48GP98	$HD^+ + H_2 \rightarrow H_2D^+ + H$	38
RC49GP98	$HD^+ + H_2 \rightarrow H_3^+ + D$	38
RC50GP98	$D + H_3^+ \rightarrow H_2D^+ + H$	3
RC51GP98	$H_2D^+ + e \rightarrow 2H + D$	8, 17
RC52GP98	$H_2D^+ + e \rightarrow H_2 + D$	8, 17
RC53GP98	$H_2D^+ + e \rightarrow HD + H$	8, 17
RC54GP98	$H_2D^+ + H_2 \rightarrow H_3^+ + HD$	2, 22, 13
RC55GP98	$H_2D^+ + H \rightarrow H_3^+ + D$	3
RC76GA08	$D + H + H \rightarrow HD + H$	9
RC76Inv	$HD + H \rightarrow D + H + H$	9
RC77GA08	$D + H + HD \rightarrow HD + HD$	9
RC77Inv	$HD + HD \rightarrow D + H + HD$	9

References. — (1) Abel et al. (1997); (2) Adams & Smith (1981); (3) Adams & Smith (1985); (4) Aldrovandi & Pequignot (1973); (5) AMDIS (1989); (6) Black (1979); (7) Dalgarno & Lepp (1987); (8) Datz et al. (1995); (9) Flower & Harris (2007) ;(10) Galli & Palla (1998); (11) Gerlich (1982); (12) Gerlich & Horning (1992); (13) Herbst (1982); (14) Janev et al. (1987); (15) Karpas et al. (1979); (16) Kimura et al. (1993); (17) Larsson et al. (1996); (18) Launay et al. (1991); (19) Lepp & Shull (1984); (20) MacLow & Shull (1986); (21) Mielke et al. (1994); (22) Millar et al. (1989); (23) Orient (1977); (24) Peart & Hayton (1994). (25) Peterson et al. (1971); (26) Poulaert et al. (1978); (27) Roberge & Dalgarno (1982); (28) Savin (2002); (29) Savin et al. (2004); (30) Schneider et al. (1994); (31) Schulz & Asundi (1967); (32) Shavitt (1959); (33) Sidhu et al. (1992); (34) Stancil et al. (1993); (35) Stibbe & Tennyson (1999); (36) Stromholm et al. (1995); (37) Sundstrom et al. (1994); (38) Theard & Huntress (1974); (39) Verner & Ferland (1996); (40) Wishar (1979); (41) Yousif & Mitchell (1989); (42) Zigelman et al. (1989).

TABLE 3
Rate coefficents for chemical reactions.

Reaction rate (cm^3/s)	reaction	ref
Reactions with Li		
RC56GP98	$Li^+ + e \rightarrow Li + \gamma$	8
RC57GP98	$Li^+ + H^- \rightarrow Li + H$	4
RC58GP98	$Li^- + H^+ \rightarrow Li + H$	4
RC59GP98	$Li + e \rightarrow Li^- + \gamma$	5
RC60GP98	$Li + H^+ \rightarrow Li^+ + H$	2
RC61GP98	$Li + H^+ \rightarrow Li^+ + H + \gamma$	6
RC62GP98	$Li + H^- \rightarrow LiH + e$	7
RC63GP98	$Li^- + H \rightarrow LiH + e$	7
RC64GP98	$LiH^+ + H \rightarrow LiH + H^+$	7
RC65GP98	$LiH + H^+ \rightarrow LiH^+ + H$	7
RC66GP98	$LiH + H \rightarrow Li + H_2$	7
RC67GP98	$Li^+ + H \rightarrow LiH^+ + \gamma$	1
RC68GP98	$Li + H^+ \rightarrow LiH^+ + \gamma$	1
RC69GP98	$LiH + H^+ \rightarrow LiH^+ + H$	7
RC70GP98	$LiH + H^+ \rightarrow Li^+ + H_2$	7
RC71GP98	$LiH^+ + e \rightarrow Li + H$	7
RC72GP98	$LiH^+ + H \rightarrow Li + H_2^+$	7
RC73GP98	$LiH^+ + H \rightarrow Li^+ + H_2$	7
RC78M05	$Li + H_2 \rightarrow LiH + H$	3
RC79M05	$Li + H + H \rightarrow LiH + H$	3
RC80M05	$Li + H + H_2 \rightarrow LiH + H_2$	3

References. — (1) Dalgarno et al. (1996); (2) Kimura et al. (1994); (3) Mizusawa et al. (2005); (4) Peart & Hayton (1994); (5) Ramsbottom et al. (1994); (6) Stancil & Zigelman (1996); (7) Stancil et al. (1996); (8) Verner & Ferland (1996).

TABLE 4
Cooling processess.

Cooling process (erg/cm^3s)	reaction	ref
Collisional ionization		
H		
CI1A	$H + e \rightarrow H^+ + 2e$	1
CI2A	$H^- + e \rightarrow H + 2e$	1
CI3A	$H^- + H \rightarrow 2H + e$	1
He		
CI4A	$He + e \rightarrow He^+ + e$	1
CI5A	$He^+ + e \rightarrow He^{++} + 2e$	1
D		
CI6A	$D + e \rightarrow D^+ + 2e$	1
Recombination		
H		
Rec1A97	$H^+ + e \rightarrow H + \gamma$	1,2
He		
Rec4GP98r, Rec4A97d	$He^+ + e \rightarrow He + \gamma$	1,2,3
Rec5GP98d	$He^{++} + e \rightarrow He^+ + \gamma$	3
D		
Rec9GP98	$D^+ + e \rightarrow D + \gamma$	3
Collisional excitation of		
CE1A97	H	2,1
CE2GP98	H_2	3
CE3A97	He	2,1
CE4A97	He^+	2,1
CE5GP98	HD	3
CE6GP98	LiH	3
Bremsstrahlung		2
Compton cooling		4

References. — (1) Abel et al. (1997); (2) Black (1981); (3) Galli & Palla (1998); (4) Peebles (1971); (5) Verner & Ferland (1996).

TABLE 5

Cross sections (σ_A), ionization rates (IR_A) and heatings (Γ_A).

$\sigma_A(cm^2)$ $IR_A(s^{-1})$ $\Gamma_A(erg/cm^{-3}s)$	reaction	ref
H		
σ_H1Gl07		
IR_H1Gl07		
Γ_H1Gl07	$H+\gamma \rightarrow H^+ + e$	8
σ_H2Gl07		
IR_H2Gl07		
Γ_H2Gl07	$H^- + \gamma \rightarrow H + e$	2
σ_H3Gl07		
IR_H3Gl07		
Γ_H3Gl07	$H_2 + \gamma \rightarrow H_2^+ + e$	6
IR_H3Gl07p		
Γ_H3Gl07p	$H_2 + \gamma \rightarrow H_2^* \rightarrow H + H$	4
σ_HGl407		
IR_H4Gl07		
Γ_H4Gl07	$H_2^+ + \gamma \rightarrow H^+ + H$	3
σ_H5A97		
IR_H5A97		
Γ_H5A97	$H_2^+ + \gamma \rightarrow 2H^+ + e$	11
σ_H6A97		
IR_H6A97		
Γ_H6A97	$H_2 + \gamma \rightarrow H + H$	1
He		
σ_{He}7Gl07		
IR_{He}7Gl07		
Γ_{He}7Gl07	$He + \gamma \rightarrow He^+ + e$	12
σ_{He}8A97		
IR_{He}8A97		
Γ_{He}8A97	$He^+ + \gamma \rightarrow He^{++} + e$	7
D		
σ_D9Gl07		
IR_D9Gl07		
Γ_D9Gl07	$D + \gamma \rightarrow D^+ + e$	8
σ_D10A97		
IR_D10Gl07		
Γ_D10A97	$HD^+ + \gamma \rightarrow H + D^+$	11
σ_D11A97		
IR_D11Gl07		
Γ_D11A97	$HD^+ + \gamma \rightarrow H^+ + D$	11
Li		
σ_{Li}13RM		
IR_{Li}13RM		
Γ_{Li}13RM	$Li + \gamma \rightarrow Li^+ + e$	10
σ_{Li}15Rams		
IR_{Li}15Rams		
Γ_{Li}15Rams	$Li^- + \gamma \rightarrow Li + e$	9
σ_{Li}16KD		
IR_{Li}16KD		
Γ_{Li}16KD	$LiH + \gamma \rightarrow Li + H$	5

References. — (1) Abel et al. (1997); (2) de Jong (1972); (3) Dunn (1968); (4) Galli & Palla (1998); (5) Kirby & Dalgarno (1978); (6) O'Neill & Reinhardt (1978); (7) Osterbrock (1977); (8) Osterbrock (1989); (9) Ramsbottom et al. (1994); (10) Reilman & Manson (1979); (11) Shapiro & Kang (1987); (12) Yan et al. (1998).